

ACETAMINOPHEN (TYLENOL) DOSING CHART

		Liquid Suspension or Elixir	Chewable Tablets	Junior Strength (Tablets or Capsules)	Regular Strength (Tablets or Capsules)	Extra Strength (Tablets or Capsules)
Weight (pounds)	Dose	160 mg/5 mL (teaspoon)	80 mg each	160 mg each	325 mg each	500 mg each
12 - 17	80 mg	2.5 mL (½ teaspoon)	1 tablet			
18 - 23	120 mg	3.75 mL (¾ teaspoon)	1 ½ tablets			
24 - 35	160 mg	5 mL (1 teaspoon)	2 tablets	1 tablet or capsule		
36 - 47	240 mg	7.5 mL (1 ½ teaspoons)	3 tablets	1 ½ tablets or capsules		
48 - 59	320 mg	10 mL (2 teaspoons)	4 tablets	2 tablets or capsules	1 tablet or capsule	
60 - 71	400 mg	12.5 mL (2 ½ teaspoons)	5 tablets	2 ½ tablets or capsules		
72 - 95	480 mg	15 mL (3 teaspoons)	6 tablets	3 tablets or capsules		1 tablet or capsule
> 95	650 mg	20 mL (4 teaspoons)	8 tablets	4 tablets or capsules	2 tablets or capsules	

**** ACETAMINOPHEN SHOULD NOT BE GIVEN TO AN INFANT LESS THAN 3 MONTHS OLD WITHOUT CONSULTING YOUR PHYSICIAN! ****

**** ACETAMINOPHEN MAY BE GIVEN AT INTERVALS OF EVERY 4-6 HOURS ****

IBUPROFEN (MOTRIN/ADVIL) DOSING CHART

		Infants' Oral Drops	Liquid Suspension or Elixir	Junior Strength (Chewable Tablets or Caplets)	Regular Strength (Tablets or Capsules)
Weight (pounds)	Dose	50 mg/1.25 mL	100 mg/5 mL	100 mg each	200 mg each
12 - 17	50 mg	1.25 ml	2.5 mL (½ teaspoon)		
18 - 21	75 mg	1.875 ml	3.75 mL (¾ teaspoon)		
22 - 32	100 mg	2.5 ml	5 mL (1 teaspoon)	1 chewable tablet or caplet	
33 - 43	150 mg	3.75 ml	7.5 mL (1 ½ teaspoons)	1 ½ chewable tablets or caplets	
44 - 54	200 mg	5 ml	10 mL (2 teaspoons)	2 chewable tablets or caplets	1 tablet or capsule
55 - 65	250 mg		12.5 mL (2 ½ teaspoons)	2 ½ chewable tablets or caplets	
66 - 76	300 mg		15 mL (3 teaspoons)	3 chewable tablets or caplets	
77 - 87	350 mg		17.5 mL (3 ½ teaspoons)	3 ½ chewable tablets or caplets	
88 - 132	400 mg		20 mL (4 teaspoons)	4 chewable tablets or caplets	2 tablets or capsules
> 132	600 mg		30 mL (6 teaspoons)	6 chewable tablets or caplets	3 tablets or capsules

****IBUPROFEN SHOULD NOT BE ADMINISTERED TO AN INFANT UNDER 6 MONTHS OF AGE!****

****IBUPROFEN MAY BE GIVEN AT INTERVALS OF EVERY 6-8 HOURS****